THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION

535 NORTH DEARBORN STREET - CHICAGO 10, ILL.

Cable Address - . . "Medic, Chicago"

Subscription price - - . : Eight dollars per annum in advance

Please send in promptly notice of change of address, giving both old and new; always state whether the change is temporary or permanent. Such notice should mention all journals received from this office. Important information regarding contributions will be found on second advertising page following reading matter.

SATURDAY, NOVEMBER 25, 1944

ENVIRONMENTAL CANCER

Environmental cancer is cancer due to prolonged contact with some cancerigenic agent in the environment. The relationship of a previous exposure to such an agent and the cancer often appearing many years later is frequently overlooked or not understood. Environmental cancers can be divided into four main groups according to the type of exposure:

First are the cancers caused by agents in the normal environment. Representative of this group is cancer due to the continued ingestion of arsenic in drinking water and foodstuffs in certain regions, the solar cancers of the skin in dry and sunny regions, as in our Midwestern and Southwestern states, the bladder cancers in people in places like Egypt with endemic schistosomiasis, and possibly also lung cancers due to inhalation of road dust containing cancerigenic tar. The second group of environmental cancer results from certain habits (habitual cancer), as for example cancer of the lip and mouth in smokers, cancer of the oral lining in betel and tobacco chewers, and cancer of the abdominal skin in kangri users. The third group comprises cancer resulting from use of certain medicinal agents (medicinal cancers), e. g. cancer of the skin after arsenical medication and after exposure to roeutgen rays and radioactive substances. The fourth, which is the largest and most important group, is represented by occupational cancer elicited by exposure to chemical and physical agents in the course of regular occupations.1

The agents known or suspected to cause occupational cancer are arsenic chromates, nickel carbonyl, radium, mesotherium, asbestos crude and processed mineral oils,

pitch, tar. soot, paraffin oil, anthracene oil, creasone aromatic amino compounds (aniline, napithylamine benzidine), benzene, ultraviolet rays, roention rays, radioactive materials and substances from certain parasitic worms. Approximately 8,000 to 9,000 cases of occupational cancer appear to have been claced on record, the majority during the last two to three decades, during which the sources of occupational cancer were mostly discovered. These figures, however, do not reflect the actual incidence of industrial cancer, as the cause of many cases of this type is not properly recognized or their occurrence is not recorded.

Environmental cancer may involve many organs and tissues, notably the skin, lip, tongue, cheek, oral cavity, nasal sinuses, larynx, bronchi, lung, liver, kidney, ureter, bladder, connective tissue, bone and hemopoietic tissues. The site of the cancer depends on the type of the cancerigenic agent and the kind of exposure. The sex ratio is determined mainly by the exposure to the various cancerigenic agents. It is for this reason that the male sex is predominantly affected by occupational cancer, but osteogenic sarcoma occurs in women working in luminous dial factories. Undoubtedly the rapid increase in the employment of women in hazardom industries will be followed by an increase of occupational cancer in women.

The public health importance of industrial cancer undoubtedly will increase not only with respect to the number of workers, male and female, exposed to hairstrial cancerigenic agents but also with respect to the variety of plants and operations in question. Figure health authorities will wish to institute effective ; and sanitary supervision of such establishments as real as medical control of the workers, present as according past. Practitioners and cancer clinics, in a large to industrial physicians, will aid greatly in meeting this situation by close study of the occupational for any itall patients coming to them with precance- or and cancerous lesions. It is mainly in this way if it the industrial origin of cancer can be established and new sources of occupational forms discovered. The sevents is from such efforts will be not only the early street at and improved prognosis of cases of occupations but also the prevention of industrial cancer eli nination of exposures to cancerigenic agenttrial cancer is largely preventable by proper telling mirasures.

cially occupational caucers is a promising appropriate the study of fundamental problems of human caucer because these cancers are the counterpart of expensional cancer in animals. Environmental cancers are mit a scientific analysis of cancerigenic agents and of local and general reactions induced by them in human beings.

bet of

tor

wh

οf

per

wer

tub

tati

in t

ate

bre

we

of

ext res cul lat:

Lo sio likcas Lo

 $C \in \Gamma$

the cor Ti

Dei

10 <u>đ</u>e

Oi Wi

the in:

М А.

3. ;

de Na

^{1.} Clemensen, J.: Canter and Geographical in Denmark, 1925/1919. Copenhagen, Nyr Nordisk Forlag Arnold Busck, 1941. Davis, E.: Cheminal Carcinogenesis, Drugs, Dyes, Remedies and Cosmetics with Particular Petersence to Bladder Tumors, J. Urol. 49: 14, 1943. Gross, E.: Dos Carcinom vom Standbenkt des Gewerbefoxikologen, Ztschr. f. angew. Chem. 33: 358, 1940. Hueper, W. C.: Occupational Lamors and Killed Biteases, Springhelt, Ill., Charles C. Thomas, 1942, p. 396; Canter in the Kelaron in Occupation and Environment, Bull. Am. Soc. Control of London 25: 6, 1943. Warren, S.: Minimal Criteria Required to Prove Causation of Traumatic or Occupational Neoplasms, Ann. Surg. 117: 585, 1943.